COMPUTER SYSTEM
100

MEMORY 112

COMPUTATIONAL
CODE FOR
INTERVALS
114

INTERVAL VALUES
115

PROCESSOR
102

ARITHMETIC
UNIT
104

BRIDGE
106

PERIPHERAL BUS
110

STORAGE
DEVICE
108

FIG. 1

SOURCE CODE
FOR INTERVAL
COMPUTATIONS
202

COMPILER FOR
INTERVAL CODE
204

EXECUTABLE
CODE FOR
INTERVAL
COMPUTATIONS
206

PROCESSOR
102

INTERVAL
VALUES
115

ARITHMETIC
UNIT FOR
INTERVAL
COMPUTATIONS
104

RESULTS
212

FIG. 2

INTERVAL
302

FIRST FLOATING
POINT NUMBER 304

SECOND FLOATING
POINT NUMBER 306

INTERVAL
312

FIRST FLOATING
POINT NUMBER 314

SECOND FLOATING
POINT NUMBER 316

ARITHMETIC UNIT FOR
INTERVAL
COMPUTATIONS
104

FIRST FLOATING
POINT NUMBER 324

SECOND FLOATING
POINT NUMBER 326

INTERVAL
322

FIG. 3

START
400

RECEIVE REPRESENTATION
OF INTERVAL
402

PERFORM ARITHMETIC
OPERATION USING
REPRESENTATION OF
INTERVAL 404

END
406

FIG. 4


$$X \equiv [\underline{x}, \overline{x}] = \{x \in \mathfrak{R}^* \mid \underline{x} \leq x \leq \overline{x}\}$$

$$Y \equiv [\underline{y}, \overline{y}] = \{y \in \mathfrak{R}^* \mid \underline{y} \leq y \leq \overline{y}\}$$

(1) $\quad X + Y = \left[\downarrow \underline{x} + y, \uparrow \overline{x} + \overline{y}\right]$

(2) $\quad X - Y = \left[\downarrow \underline{x} - \overline{y}, \uparrow \overline{x} - \underline{y}\right]$

(3) $\quad X \times Y = \left[\min\left(\downarrow \underline{x} \times y, \underline{x} \times \overline{y}, \overline{x} \times y, \overline{x} \times \overline{y}\right), \max\left(\uparrow \underline{x} \times y, \underline{x} \times \overline{y}, \overline{x} \times y, \overline{x} \times \overline{y}\right)\right]$

(4) $\quad X / Y = \left[\min\left(\downarrow \underline{x} / y, \underline{x} / \overline{y}, \overline{x} / y, \overline{x} / \overline{y}\right), \max\left(\uparrow \underline{x} / y, \underline{x} / \overline{y}, \overline{x} / y, \overline{x} / \overline{y}\right)\right], \text{if } 0 \notin Y$

$$X / Y \subseteq \mathfrak{R}^*, \text{if } 0 \in Y$$

FIG. 5

FIG. 6

START

FOR EACH BOX IN LIST L_1, APPLY TERM CONSISTENCY TO EACH OF THE INEQUALITY CONSTRAINTS $p_i(\mathbf{x}) \leq 0$ $(i=1, ..., m)$.
701

IF $f\_bar < +\infty$, THEN FOR EACH BOX IN L_1, APPLY TERM CONSISTENCY TO THE INEQUALITY $f \leq f\_bar$.
702

IF L_1 IS EMPTY, GO TO STEP 742. OTHERWISE, SELECT THE BOX IN L_1 FOR WHICH THE LOWER BOUND OF $f(\mathbf{X})$ IS SMALLEST. FOR LATER REFERENCE, DENOTE THIS BOX BY $\mathbf{X}^{(1)}$. DELETE $\mathbf{X}$ FROM L_1.
703

APPLY TERM CONSISTENCY OVER $\mathbf{X}$ TO EACH CONSTRAINT INEQUALITY. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
704

COMPUTE AN APPROXIMATION x FOR THE CENTER $m(\mathbf{X})$ OF $\mathbf{X}$. IF f(x) > f_bar, GO TO STEP 708.
705

FOR FUTURE REFERENCE, DENOTE THE BOX $\mathbf{X}$ by $\mathbf{X}^{(2)}$. DO A CONSTRAINED LINE SEARCH TO TRY TO REDUCE f_bar.
706

IF f_bar WAS NOT REDUCED IN STEP 706, GO TO STEP 709.
707

APPLY TERM CONSISTENCY TO THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$ OVER THE CURRENT BOX $\mathbf{X}$. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
708

IF $w(\mathbf{X}) < \varepsilon_X$ AND $w[f(\mathbf{X})] < \varepsilon_F$, PUT $\mathbf{X}$ IN LIST L_2. OTHERWISE, IF $\mathbf{X}$ IS SUFFICIENTLY REDUCED RELATIVE TO THE BOX $\mathbf{X}^{(1)}$, PUT $\mathbf{X}$ IN L_1 AND GO TO STEP 703.
709

APPLY BOX CONSISTENCY TO EACH INEQUALITY CONSTRAINT. IF $f\_bar < +\infty$, APPLY BOX CONSISTENCY TO THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
710

IF THE UPPER BOUND OF $p_i(\mathbf{X}) \geq 0$ FOR ANY $i=1, ..., n$, GO TO STEP 726.
711

APPLY TERM CONSISTENCY TO $g_i=0$ FOR $i=1, ..., n$. IF THE RESULT FOR ANY $i=1, ..., n$ IS EMPTY, GO TO STEP 703.
712

A

FIG. 7A

(A)

APPLY TERM CONSISTENCY TO THE RELATION $H_{ii}(\mathbf{x}) \geq 0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
713

REPEAT STEP 709.
714

APPLY BOX CONSISTENCY TO $g_i=0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
715

APPLY BOX CONSISTENCY TO $H_{ii}(\mathbf{x}) \geq 0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
716

REPEAT STEP 709.
717

USE A CRITERION $w(\mathbf{X}) > (w_I + w_R)/2$ TO DECIDE IF A NEWTON STEP SHOULD BE APPLIED TO SOLVE $\mathbf{g}=\mathbf{0}$. IF NOT, GO TO STEP 726.
718

GENERATE THE INTERVAL JACOBIAN $\mathbf{J}(\mathbf{x},\mathbf{X})$ OF THE GRADIENT $\mathbf{g}$ AND COMPUTE THE APPROXIMATE INVERSE $\mathbf{B}$ OF THE CENTER OF $\mathbf{J}(\mathbf{x},\mathbf{X})$. APPLY ONE STEP OF AN INTERVAL NEWTON METHOD TO SOLVE $\mathbf{g}=\mathbf{0}$. IF THE RESULT IS EMPTY, GO TO STEP 703.
719

REPEAT STEP 709.
720

OBTAIN $\mathbf{Bg}$ IN ANALYTIC FORM. APPLY TERM CONSISTENCY TO SOLVE THE i^{th} EQUATION OF $\mathbf{Bg}=\mathbf{0}$ FOR THE i^{th} VARIABLE x_i FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
721

REPEAT STEP 709.
722

USE BOX CONSISTENCY TO SOLVE THE i^{th} EQUATION OF $\mathbf{Bg}$ FOR THE i^{th} VARIABLE FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
723

REPEAT STEP 709.
724

(B)

FIG. 7B

(B)

USE THE MATRIX B IN A NEWTON STEP TO TRY TO REDUCE *f_bar*.
725

COMPUTE AN APPROXIMATION x FOR THE CENTER $m(\mathbf{X})$ OF **X**. IF $f(\mathbf{x}) > f\_bar$, GO TO STEP 703.
726

SKIP THIS STEP AND GO TO STEP 732 IF $\mathbf{X} = \mathbf{X}^{(2)}$ IS THE SAME BOX FOR WHICH A LINE SEARCH WAS DONE IN STEP 706. OTHERWISE, DO A LINE SEARCH TO TRY TO REDUCE *f_bar*. IF *f_bar* IS NOT REDUCED, GO TO STEP 732.
727

FOR FUTURE REFERENCE DENOTE $\mathbf{X}^{(3)} = \mathbf{X}$. USE A LINEARIZATION TEST TO DECIDE WHETHER TO LINEARIZE AND "SOLVE" THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$. IF THE CRITERION IS NOT SATISFIED, GO TO STEP 732.
728

USE A LINEAR METHOD TO TRY TO REDUCE **X** USING THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$. IF **X** IS DELETED, GO TO STEP 703. OTHERWISE, IF THIS APPLICATION OF THE LINEAR METHOD DOES NOT SUFFICIENTLY REDUCE BOX $\mathbf{X}^{(3)}$ GO TO STEP 731.
729

USE A QUADRATIC METHOD TO TRY TO REDUCE **X** USING THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$. IF **X** IS DELETED, GO TO STEP 703.
730

REPEAT STEP 709.
731

USE A LINEARIZATION TEST TO DECIDE WHETHER TO LINEARIZE AND "SOLVE" THE INEQUALITY CONSTRAINTS. IF THE PROCEDURE INDICATES THAT THE LINEARIZATION SHOULD NOT BE DONE, GO TO STEP 739.
732

SELECT THE INEQUALITY CONSTRAINTS TO BE SOLVED IN LINEARIZED FORM, AND POSSIBLY ADD TO THIS SET THE INEQUALITY $f(\mathbf{x}) \leq f\_bar$. IF NO INEQUALITIES ARE SELECTED, GO TO STEP 739. OTHERWISE, LINEARIZE THE RESULTING SET OF INEQUALITIES, AND SOLVE THE RESULTING SET OF LINEAR INEQUALITIES. IF THE SOLUTION SET IS EMPTY, GO TO STEP 703.
733

REPEAT STEP 709.
734

(C)

FIG. 7C

(C)

USE PRECONDITIONING MATRIX **B** TO ANALYTICALLY PRECONDITION THE SET OF INEQUALITIES THAT WERE SELECTED FOR USE IN STEP 733. USE TERM CONSISTENCY TO SOLVE EACH PRECONDITIONED INEQUALITY.
735

REPEAT STEP 709.
736

USE BOX CONSISTENCY TO SOLVE THE SAME INEQUALITIES FOR THE SAME VARIABLES AS IN STEP 735.
737

REPEAT STEP 709.
738

USE A LINEARIZATION TEST TO DECIDE WHETHER TO SOLVE THE JOHN CONDITIONS. IF NOT, GO TO STEP 742.
739

MODIFY THE JOHN CONDITIONS BY OMITTING THOSE CONSTRAINTS p_i FOR WHICH $p_i(\mathbf{X}) < 0$ (SINCE THEY ARE NOT ACTIVE IN **X**). APPLY ONE PASS OF THE INTERVAL NEWTON METHOD TO THE (MODIFIED) JOHN CONDITIONS. IF THE RESULT IS EMPTY, GO TO STEP 703.
740

REPEAT STEP 709.
741

IN VARIOUS PREVIOUS STEPS, GAPS MAY HAVE BEEN FORMED IN COMPONENTS OF **X**. IF SO, MERGE ANY OVERLAPPING GAPS. SPLIT **X**, AND PLACE THE RESULTING SUBBOXES IN L_1 AND GO TO STEP 703.
742

IF $f\_bar < +\infty$, APPLY TERM CONSISTENCY TO $f(\mathbf{x}) \leq f\_bar$ FOR EACH BOX IN THE LIST L_2. DENOTE THOSE THAT REMAIN BY $\mathbf{X}^{(1)}, ..., \mathbf{X}^{(s)}$ WHERE s IS THE NUMBER OF BOXES REMAINING. DETERMINE

$$\underline{F} = \min_{1 \leq i \leq s} \underline{f}(X^{(i)}) \text{ and } \overline{F} = \max_{1 \leq i \leq s} \overline{f}(X^{(i)})$$

743

TERMINATE.
744

FIG. 7D